arXiv:1211.0667v1 [hep-ph] 4 Nov 2012

Relativistic quark-gluon description of ^3He

S.M. Gerasyuta* and E.E. Matskevich†
Department of Theoretical Physics, St. Petersburg State University, 198904, St. Petersburg, Russia and
Department of Physics, LTA, 194021, St. Petersburg, Russia

The relativistic nine-quark equations are found in the framework of the dispersion relation technique. ^3He nucleus is described by these equations. We consider the ^3He as the system of interacting quarks and gluons. The approximate solutions of these equations using the method based on the extraction of leading singularities of the amplitudes are obtained. The relativistic nine-quark amplitudes of ^3He, including the u, d quarks are calculated. The poles of these amplitudes determine the mass of nine-quark system. The ^3He mass $M = 2809\,MeV$ is calculated. The gluon coupling constant in the light nuclei region is equal to $g = 0.1536$. The gluon interaction of ^3He is obtained in $2-3$ time smaller as compared with baryon interaction.

PACS numbers: 11.55.Fv, 12.39.Ki, 12.39.Mk, 12.40.Yx.

I. INTRODUCTION.

In nuclear physics one studies interacting hadrons in the nonperturbative regime of quantum chromodynamics (QCD). This field faces two natural frontiers. The first one is devoted to the origin and the fundaments of the strong force between the nucleous, as well as its parametrization in terms of effective degrees of freedom, suitable for the nonperturbative regime. The other is the completely microscopic, quantum mechanical treatment of many-body systems (nuclei) in terms of these effective degrees of freedom with the strong force as dynamical input. Within these two frontiers, few-nucleon physics at low energy represents quite a rich and flourishing research field describing a greate variety of strong interaction phenomena.

The study of polarization phenomena in hadron and hadron-nucleus interactions gives more detailed information on dynamics of their interactions and the structure of colliding particles. The quark structure and relativistic effects of light nuclei, in particular, deuterons, is one of the important problems in nuclear physics at low and intermediate energies. The theoretical and experimental study of reactions like the elastic $e-d$ [1] and $p-d$ [2, 3] scattering, deuteron break-up reactions induced by electrons or protons [4, 5] and the deuteron stripping processes on protons and nuclei at the low and intermediate energies [6, 7], can allow us to find out new information on the deuteron structure at short distances. The elastic backward proton-deuteron scattering has been experimentally and theoretically studied in Saclay [2], Dubna and at JLab [8–11]. Usually these processes are analyzed within a simple impulse approximation. Up to now all these data have not been described within the one-nucleon exchange model (ONE), including even the relativistic effects in the deuteron [12, 13]. Then the elastic backward proton-deuteron scattering within the relativistic approach including the ONE and the high order graphs corresponds to the emission, rescattering and absorption of the virtual pion by a nucleon of deuteron [14].

The main aim of the investigation of the reaction by the polarized deuterons is to establish the nature of $2N$ and $3N$ forces, the role of the relativistic effects and nonnucleon degrees of freedom. The last decades such investigation were performed at different experiments all over the world at RIKEN, KVI, IUCF and RCNP. The activing was stimulated by the discrepancy of 30% between the measured cross section for deuteron-proton elastic scattering at intermediate energies and the Faddeev calculations using modern potentials of nucleon-nucleon interaction.

In Ref. [15] a representation of the Faddeev equation in the form of a dispersion relation in the pair energy of the two interacting particles was used. This was found to be convenient in order to obtain an approximate solution of the Faddeev equation by a method based on extraction of leading singularities of the amplitude. With a rather crude approximation of the low-energy NN interaction a relatively good description of the form factor of tritium (helium-3) at low q^2 was obtained.

In our papers [16–18] relativistic generalization of the three-body Faddeev equations was obtained in the form of dispersion relations in the pair energy of two interacting quarks. The mass spectrum of S-wave baryons including u, d, s quarks was calculated by a method based on isolating the leading singularities in the amplitude. We searched for the approximate solution of integral three-quark equations by taking into account two-particle and triangle singularities,

---

[*] Electronic address: gerasyuta@SG6488.spb.edu
[†] Electronic address: matskev@pobox.spbu.ru

all the weaker ones being neglected. If we considered such an approximation, which corresponds to taking into account two-body and triangle singularities, and defined all the smooth functions of the subenergy variables (as compared with the singular part of the amplitude) in the middle point of the physical region of Dalitz-plot, then the problem was reduced to the one of solving a system of simple algebraic equations.

In the recent paper [19] the relativistic six-quark equations are found in the framework of coupled-channel formalism. The dynamical mixing between the subamplitudes of hexaquark are considered. The six-quark amplitudes of dibaryons are calculated. The poles of these amplitudes determine the masses of dibaryons. We calculated the contribution of six-quark subamplitudes to the hexaquark amplitudes.

In the present paper the ^3He as the system of interacting quarks and gluons is considered. The relativistic nine-quark equations are found in the framework of the dispersion relation technique. The dynamical mixing between the subamplitudes of ^3He is taken into account. The relativistic nine-quark amplitudes of ^3He, including the u, d quarks are calculated. The approximate solutions of these equations using the method based on the extraction of leading singularities of the amplitude are calculated. The poles of the nine-quark amplitudes determine the mass of ^3He. The mass of ^3He state is equal to $M = 2809\,MeV$. The model use only two parameters: cutoff $\Lambda = 11$ [19] and gluon coupling constant $g = 0.1536$, which is smaller then the gluon coupling constant of hadron physics.

In Sec. II, we briefly discuss the relativistic Faddeev approach. The relativistic three-quark equations are constructed in the form of the dispersion relation over the two-body subenergy. The approximate solution of these equations using the method based on the extraction of leading singularities of the amplitude are obtained. We calculated the mass spectrum of S-wave baryons with $J^P = \frac{1}{2}^+, \frac{3}{2}^+$ (Table I).

In Sec. III, the nine-quark amplitudes of ^3He are constructed. The dynamical mixing between the subamplitudes of ^3He is considered. The relativistic nine-quark equations are constructed in the form of the dispersion relation over the two-body subenergy. The approximate solutions of these equations using the method based on the extraction of leading singularities of the amplitude are obtained.

Sec. IV is devoted to the calculation results for the ^3He (^3H).

In conclusion, the status of the considered model is discussed.

II. BRIEF INTRODUCTION OF RELATIVISTIC FADDEEV EQUATIONS.

We consider the derivation of the relativistic generalization of the Faddeev equation for the example of the Δ-isobar ($J^P = \frac{3}{2}^+$). This is convenient because the spin-flavour part of the wave function of the Δ-isobar contains only nonstrange quarks and pair interactions with the quantum numbers of a $J^P = 1^+$ diquark (in the color state $\bar{3}_c$). The $3q$ baryon state Δ is constructed as color singlet. Taking into account the equality of all pair interactions of nonstrange quarks in the state with $J^P = 1^+$, we obtain the corresponding equation for the amplitudes (Fig. 1):

$$A_1(s, s_{12}, s_{13}, s_{23}) = \lambda + A_1(s, s_{12}) + A_1(s, s_{13}) + A_1(s, s_{23})\,. \tag{1}$$

Here, the s_{ik} are the pair energies of particles 1, 2 and 3, and s is the total energy of the system. To write down a concrete equations for the function $A_1(s, s_{12})$ we must specify the amplitude of the pair interaction of the quarks. We write the amplitude of the interaction of two quarks in the state $J^P = 1^+$ in the form:

$$a_1(s_{12}) = \frac{G_1^2(s_{12})}{1 - B_1(s_{12})}\,, \tag{2}$$

$$B_1(s_{12}) = \int\limits_{4m^2}^{\infty} \frac{ds'_{12}}{\pi} \frac{\rho_1(s'_{12}) G_1^2(s'_{12})}{s'_{12} - s_{12}}\,, \tag{3}$$

$$\rho_1(s_{12}) \;=\; \left(\frac{1}{3}\,\frac{s_{12}}{4m^2} + \frac{1}{6}\right) \left(\frac{s_{12} - 4m^2}{s_{12}}\right)^{\frac{1}{2}}\,. \tag{4}$$

Here $G_1(s_{12})$ is the vertex function of a diquark with $J^P = 1^+$. $B_1(s_{12})$ is the Chew-Mandelstam function [20], and $\rho_1(s_{12})$ is the phase spaces for a diquark with $J^P = 1^+$.

The pair quarks amplitudes $qq \to qq$ are calculated in the framework of the dispersion N/D method with the input four-fermion interaction [21–23] with the quantum numbers of the gluon [24, 25].

The four-quark interaction is considered as an input:

$$g_V\,(\bar{q}\lambda I_f\gamma_\mu q)^2+2\,g_V^{(s)}\,(\bar{q}\lambda I_f\gamma_\mu q)\,(\bar{s}\lambda\gamma_\mu s)+g_V^{(ss)}\,(\bar{s}\lambda\gamma_\mu s)^2\,. \tag{5}$$

Here I_f is the unity matrix in the flavor space (u,d). λ are the color Gell-Mann matrices. Dimensional constants of the four-fermion interaction g_V, $g_V^{(s)}$ and $g_V^{(ss)}$ are parameters of the model. At $g_V=g_V^{(s)}=g_V^{(ss)}$ the flavor $SU(3)_f$ symmetry occurs. The strange quark violates the flavor $SU(3)_f$ symmetry. In order to avoid additional violation parameters we introduce the scale of the dimensional parameters [25]:

$$g=\frac{m^2}{\pi^2}g_V=\frac{(m+m_s)^2}{4\pi^2}g_V^{(s)}=\frac{m_s^2}{\pi^2}g_V^{(ss)}\,,\qquad\qquad \Lambda=\frac{4\Lambda(ik)}{(m_i+m_k)^2}\,. \tag{6}$$

Here m_i and m_k are the quark masses in the intermediate state of the quark loop. Dimensionless parameters g and Λ are supposed to be constants which are independent of the quark interaction type.

In the case under consideration the interacting pairs of particles do not form bound states. Thefore, the integration in the dispersion integral (7) run from $4m^2$ to ∞. The equation corresponding to Fig. 1 can be written in the form:

$$\begin{aligned} A_1(s,s_{12}) \;&=\; \frac{\lambda_1 B_1(s_{12})}{1-B_1(s_{12})}+\frac{G_1(s_{12})}{1-B_1(s_{12})}\int\limits_{4m^2}^{\infty}\frac{ds'_{12}}{\pi}\frac{\rho_1(s'_{12})}{s'_{12}-s_{12}}G_1(s'_{12}) \\ &\times\int\limits_{-1}^{+1}\frac{dz}{2}[A_1(s,s'_{13})+A_1(s,s'_{23})]\,. \end{aligned} \tag{7}$$

In Eq. (7) z is the cosine of the angle between the relative momentum of particles 1 and 2 in the intermediate state and the momentum of the third particle in the final state in the c.m.s. of the particles 1 and 2. In our case of equal mass of the quarks 1, 2 and 3, s'_{13} and s'_{12} are related by the equation [26]

$$s'_{13}=2m^2-\frac{(s'_{12}+m^2-s)}{2}\pm\frac{z}{2}\sqrt{\frac{(s'_{12}-4m^2)}{s'_{12}}(s'_{12}-(\sqrt{s}+m)^2)(s'_{12}-(\sqrt{s}-m)^2)}\,. \tag{8}$$

The expression for s'_{23} is similar to (8) with the replacement $z\to -z$. This makes it possible to replace $[A_1(s,s'_{13})+A_1(s,s'_{23})]$ in (7) by $2A_1(s,s'_{13})$.

From the amplitude $A_1(s,s_{12})$ we shall extract the singularities of the diquark amplitude:

$$A_1(s,s_{12})=\frac{\alpha_1(s,s_{12})B_1(s_{12})}{1-B_1(s_{12})}\,. \tag{9}$$

The equation for the reduced amplitude $\alpha_1(s,s_{12})$ can be written as

$$\alpha_1(s,s_{12}) \;=\; \lambda+\frac{1}{B_1(s_{12})}\int\limits_{4m^2}^{\infty}\frac{ds'_{12}}{\pi}\frac{\rho_1(s'_{12})}{s'_{12}-s_{12}}G_1(s'_{12})\int\limits_{-1}^{+1}\frac{dz}{2}\frac{2\alpha_1(s,s'_{13})B_1(s'_{13})}{1-B_1(s'_{13})}\,. \tag{10}$$

The next step is to include into (10) a cutoff at large s'_{12}. This cutoff is needed to approximate the contribution of the interaction at short distances. In this connection we shall rewrite Eq. (10) as

$$\alpha_1(s,s_{12}) \;=\; \lambda+\frac{1}{B_1(s_{12})}\int\limits_{4m^2}^{\infty}\frac{ds'_{12}}{\pi}\Theta(\Lambda-s'_{12})\frac{\rho_1(s'_{12})}{s'_{12}-s_{12}}G_1\int\limits_{-1}^{+1}\frac{dz}{2}\frac{2\alpha_1(s,s'_{13})B_1(s'_{13})}{1-B_1(s'_{13})}\,. \tag{11}$$

The construction of the approximate solution of Eq. (11) is based on extraction of the leading singularities are close to the region $s_{ik}\approx 4m^2$. The structure of the singularities of amplitudes with a different number of rescattering is the following [26]. The strongest singularities in s_{ik} arise from pair rescatterings of quarks: square-root singularity corresponding to a threshold and pole singularities corresponding to bound states. We use the approximation in which the singularity corresponding to a single interaction of all three particles, the triangle singularity, is taken into account. Such a classification of singularities makes it possible to search for an approximate solution of Eq. (11), taking into account a definite number of leading singularities and neglecting the weaker ones.

For fixed values of s and s'_{12} the integration is carried out over the region of the variable s'_{13} corresponding to a physical transition of the current into three quarks (the physical region of Dalitz plot). It is convenient to take the central point of this region, corresponding to $z = 0$, to determinate the function $\alpha_1(s, s_{12})$ and also the Chew-Mandelstam function $B_1(s_{12})$ at the point $s_{12} = s_0 = \frac{s}{3} + m^2$. Then the equation for the Δ isobar takes the form

$$\alpha_1(s, s_0) = \lambda + I_{1,1}(s, s_0) \cdot 2\, \alpha_1(s, s_0)\,, \tag{12}$$

$$I_{1,1}(s, s_0) = \int\limits_{4m^2}^{\Lambda_1} \frac{ds'_{12}}{\pi} \frac{\rho_1(s'_{12})}{s'_{12} - s_{12}} G_1 \int\limits_{-1}^{+1} \frac{dz}{2} \frac{G_1}{1 - B_1(s'_{13})}\,. \tag{13}$$

We can obtain an approximate solution of Eq. (14)

$$\alpha_1(s, s_0) = \lambda [1 - 2\, I_{1,1}(s, s_0)]^{-1}\,. \tag{14}$$

The function $I_{1,1}(s, s_0)$ takes into account correctly the singularities corresponding to the fact that all propagators of triangle diagrams reduce to zero. The right-hand side of (14) may have a pole in s, which corresponds to a bound state of the three quarks. The choice of the cutoff Λ makes it possible to fix the value of the Δ isobar mass.

Baryons of S-wave multiplets have a completely symmetric spin-flavor part of the wave function, and spin $\frac{3}{2}$ corresponds to the decuplet which has a symmetric flavor part of the wave function. Octet states have spin $\frac{1}{2}$ and a mixed symmetry of the flavor function.

In analogy with the case of the Δ isobar we can obtain the rescattering amplitudes for all S-wave states with $J^P = \frac{3}{2}^+$, which include quarks of various flavors. These amplitudes will satisfy systems of integral equations. In considering the $J^P = \frac{1}{2}^+$ octet we must include the integration of the quarks in the 0^+ and 1^+ states (in the colour state $\bar{3}_c$). Including all possible rescattering of each pair of quarks and grouping the terms according to the final states of the particles, we obtain the amplitudes A_0 and A_1, which satisfy the corresponding systems of integral equations. If we choose the approximation in which two-particle and triangle singularities are taken into account, and if all functions which depend on the physical region of the Dalitz plot, the problem of solving the system of integral equations reduces to one of solving simple algebraic equations.

In our calculation the quark masses $m_u = m_d = m$ and m_s, are not uniquely determined. In order to fix m and m_s, anyhow, we make the simple assumption that $m = \frac{1}{3} m_\Delta(1232)$, $m_s = \frac{1}{3} m_\Omega(1672)$. The strange quark breaks the flavor $SU(3)_f$ symmetry (6).

In Ref. [18] we consider two versions of calculations. In the first version the $SU(3)_f$ symmetry is broken by the scale shift of the dimensional parameters. A single cutoff parameter in pair energy is introduced for all diquark states $\lambda_1 = 12.2$.

In the Table I the calculated masses of the S-wave baryons are shown. In the first version, we use only three parameters: the subenergy cutoff λ and the vertex function g_0, g_1, which corresponds to the quark-quark interaction in 0^+ and 1^+ states. In this case, the mass values of strange baryons with $J^P = \frac{1}{2}^+$ are less than the experimental ones. This means that the contribution color-magnetic is too large. In the second version, we introduce four parameters: cutoff λ_0, λ_1 and the vertex function g_0, g_1. We decrease the color-magnetic interaction in 0^+ strange channels and calculated mass values of two baryonic multiplets $J^P = \frac{1}{2}^+$, $\frac{3}{2}^+$ are in good agreement with the experimental data [27].

The essential difference between Σ and Λ is the spin of the lighter diquark. The model explains both the sign and magnitude of this mass splitting.

The suggested method of the approximate solution of the relativistic three-quark equations allows us to calculate the S-wave baryons mass spectrum. The interaction constants, determined the baryons spectrum in our model, are similar to ones in the bootstrap quark model of S-wave mesons [25]. The diquark interaction forces are defined by the gluon exchange. The relative contribution of the instanton-induced interaction is less than that with the gluon exchange. This is the consequence of $1/N_c$-expansion.

TABLE I: S-wave baryon masses $M(J^p)$ (GeV).

	$M(\frac{1}{2}^+)$			$M(\frac{3}{2}^+)$	
N	0.940	(0.940)	Δ	1.232	(1.232)
	0.940			1.232	
Λ	1.022	(1.116)	Σ^*	1.377	(1.385)
	1.098			1.377	
Σ	1.050	(1.193)	Ξ^*	1.524	(1.530)
	1.193			1.524	
Ξ	1.162	(1.315)	Ω	1.672	(1.672)
	1.325			1.672	

The gluon exchange corresponds to the color-magnetic interaction, which is responsible for the spin-spin splitting in the hadron models. The sign of the color-magnetic term is such as to made any baryon of spin $\frac{3}{2}$ heavier than its spin-$\frac{1}{2}$ counterpart (containing the same flavors).

III. NINE-QUARK AMPLITUDES OF ^3He (^3H).

We derive the relativistic nine-quark equations in the framework of the dispersion relation technique. We use only planar diagrams; the other diagrams due to the rules of $1/N_c$ expansion [28–30] are neglected. The current generates a nine-quark system. The correct equations for the amplitude are obtained by taking into account all possible subamplitudes. It corresponds to the division of complete system into subsystems with a smaller number of particles. Then one should represent a nine-particle amplitude as a sum of 36 subamplitudes:

$$A = \sum_{\substack{i<j \\ i,j=1}}^{9} A_{ij} \,. \tag{15}$$

This defines the division of the diagrams into groups according to the certain pair interaction of particles. The total amplitude can be represented graphically as a sum of diagrams. We need to consider only one group of diagrams and the amplitude corresponding to them, for example A_{12}. We shall consider the derivation of the relativistic generalization of the Faddeev-Yakubovsky approach.

In our case, the *ppn* and *pnn* states (^3He and ^3H) are considered. The isospin of ^3He is equal to $I = \frac{1}{2}$ and the spin-parity $J^P = \frac{1}{2}^+$. We take into account the pairwise interaction of all nine quarks in the nonaquark. The set of diagrams associated with the amplitude A_{12} can further be broken down into 15 groups corresponding to subamplitudes. For instance, we consider $A_1^{1^{uu}}(s, s_{12345678}, s_{1234567}, s_{123456}, s_{12345}, s_{1234}, s_{123}, s_{12})$. Here s_{ik} is the two-particle subenergy squared, s_{ijk} corresponds to the energy squared of particles i, j, k, $s_{ijk\ldots}$ is the energy squared of particles i, j, k, … and s is the system total energy squared.

In order to represent the subamplitudes A_i in the form of a dispersion relation, it is necessary to define the amplitude of qq interactions. It is similar to the three-quark case (Sec. II). We use the results of our relativistic quark model [23]

$$a_n(s_{ik}) = \frac{G_n^2(s_{ik})}{1 - B_n(s_{ik})} \,, \tag{16}$$

$$B_n(s_{ik}) = \int\limits_{4m^2}^{\Lambda} \frac{ds'_{ik}}{\pi} \frac{\rho_n(s'_{ik}) G_n^2(s'_{ik})}{s'_{ik} - s_{ik}} \,. \tag{17}$$

Here $G_n(s_{ik})$ are the diquark vertex functions. The vertex functions are determined by the contribution of the crossing channels. The vertex functions satisfy the Fierz relations. These vertex functions are generated from g_V. $B_n(s_{ik})$ and $\rho_n(s_{ik})$ are the Chew-Mandelstam functions with cutoff Λ [24] and the phase spaces:

TABLE II: Vertex functions and Chew-Mandelstam coefficients.

n	J^P	$G_n^2(s_{kl})$	α_n	β_n
1	0^+	$\frac{4g}{3} - \frac{8gm^2}{(3s_{kl})}$	$\frac{1}{2}$	0
2	1^+	$\frac{2g}{3}$	$\frac{1}{3}$	$\frac{1}{6}$

$$\rho_n(s_{ik}, J^P) = \left(\alpha(n, J^P)\frac{s_{ik}}{4m^2} + \beta(n, J^P)\right)\sqrt{\frac{s_{ik} - 4m^2}{s_{ik}}}\,. \tag{18}$$

The coefficients $\alpha(n, J^P)$, $\beta(n, J^P)$ and diquark vertex functions are given in Table II. Here $n = 1$ coresponds to qq-pairs with $J^P = 0^+$, $n = 2$ corresponds to the qq pairs with $J^P = 1^+$. In the case in question the interacting quarks do not produce a bound states, thefore the integration in Eqs. (17) is carried out from the threshold $4m^2$ to the cutoff Λ.

Let us extract singularities in the coupled equations and obtain the reduced amplitudes α_i.

The reduced amplitudes α_1 are determined by the channels 1^{uu}, 0^{ud}, 1^{dd}, the α_2 are constructed as $1^{uu}1^{uu}$, $1^{uu}0^{ud}$, $0^{ud}0^{ud}$, $1^{uu}1^{dd}$, $1^{dd}0^{ud}$, $1^{dd}1^{dd}$ channels. The reduced amplitudes α_3 are just the ppn states: $1^{uu}1^{uu}0^{ud}$, $1^{uu}1^{uu}1^{dd}$, $1^{uu}0^{ud}0^{ud}$, $1^{uu}0^{ud}1^{dd}$, $0^{ud}0^{ud}0^{ud}$, $0^{ud}0^{ud}1^{dd}$.

We consider the 15 coupled equations. The contributions of two diquarks and other 5 quarks mix with the pp and pn +3 quarks contributions. The α_3 do not include the $\alpha_3^{1^{uu}1^{dd}1^{dd}}$ state.

We used the classification of singularities, which was proposed in paper [26]. The construction of the approximate solution of equation for the α_i is based on extraction of the leading singularities of the amplitudes. The main singularities in $s_{ik} = 4m^2$ are from pair rescattering of the particles i and k. First of all there are threshold square-root singularities. Also possible are pole singularities which correspond to the bound states. We take into account more weaker singularities.

In the Fig. 2 the reduced amplitude $\alpha_1^{1^{uu}}$ is given.

The system of graphical equations in the Fig. 2 determines by the subamplitudes using the self-consistent method. The coefficients are determined by the permutation of quarks [31, 32]. We should use the coefficients multiplying of the diagrams in the graphical equation Fig. 2.

In the Fig. 2 the first coefficient is equal to 6, that the number $6 = 2$ (permutation particles 1 and 2) $\times 3$ (we can use third, 4-th, 5-th u-quarks); the second coefficient equal to 8, that the number $8 = 2$ (permutation particles 1 and 2) $\times 4$ (we can use third, 4-th, 5-th, 6-th d-quarks); the third coefficient equal to 6: $6 = 3$ (we can replace third u-quark with any of two remained u-quarks) $\times 2$ (then replace 4-th u-quark with remained u-quark); the 4-th coefficient equal to 24: $24 = 2$ (we can replace third u-quark with 4-th d-quark) $\times 3$ (we can replace third u-quark with any of two remained u-quarks) $\times 4$ (we can replace 4-th d-quark with any of three remained d-quarks); the 5-th coefficient equal to 12: $12 = 4$ (we can replace third d-quark with any of three remained d-quarks) $\times 3$ (we can replace 4-th d-quark with any of two remained d-quarks).

The similar approach allows us to take into account the coefficients in all equations.

With this classification, one defines the reduced amplitudes α_1, α_2, α_3 as well as the B-functions in the middle point of physical region of Dalitz-plot at the point s_0:

$$s_0 = \frac{s + 63}{36}\,, \quad s_{ijk} = 3s_0 - 3\,, \quad s_{ijkl} = 6s_0 - 8\,, \tag{19}$$

$$s = \sum_{\substack{i,j=1\\ i<j}}^{9} s_{ij} - 63m^2\,, \quad s_{ij} = s_0 m^2\,, \quad s = \left(\frac{M}{m}\right)^2\,. \tag{20}$$

Such choice of point s_0 allows us to replace integral equations of Fig. 2 by the algebraic equations, for instance:

$$\begin{aligned}\alpha_1^{1^{uu}} &= \lambda + 6\,\alpha_1^{1^{uu}} I_1(1^{uu}1^{uu}) + 8\,\alpha_1^{0^{ud}} I_1(1^{uu}0^{ud}) + 6\,\alpha_2^{1^{uu}1^{uu}} I_2(1^{uu}1^{uu}1^{uu}) + 24\,\alpha_2^{1^{uu}0^{ud}} I_2(1^{uu}1^{uu}0^{ud}) \\ &+\ 12\,\alpha_2^{0^{ud}0^{ud}} I_2(1^{uu}0^{ud}0^{ud})\,. \end{aligned} \tag{21}$$

$$I_1(ij) = \frac{B_j(s_0^{13})}{B_i(s_0^{12})} \int\limits_{4m^2}^{\Lambda} \frac{ds'_{12}}{\pi} \frac{G_i^2(s_0^{12})\rho_i(s'_{12})}{s'_{12}-s_0^{12}} \int\limits_{-1}^{+1} \frac{dz_1(1)}{2} \frac{1}{1-B_j(s'_{13})}\,, \tag{22}$$

$$\begin{aligned} I_2(ijk) &= \frac{B_j(s_0^{13})B_k(s_0^{24})}{B_i(s_0^{12})} \int\limits_{4m^2}^{\Lambda} \frac{ds'_{12}}{\pi} \frac{G_i^2(s_0^{12})\rho_i(s'_{12})}{s'_{12}-s_0^{12}} \frac{1}{2\pi} \int\limits_{-1}^{+1} \frac{dz_1(2)}{2} \int\limits_{-1}^{+1} \frac{dz_2(2)}{2} \\ &\times \int\limits_{z_3(2)^-}^{z_3(2)^+} dz_3(2) \frac{1}{\sqrt{1-z_1^2(2)-z_2^2(2)-z_3^2(2)+2z_1(2)z_2(2)z_3(2)}} \\ &\times \frac{1}{1-B_j(s'_{13})} \frac{1}{1-B_k(s'_{24})}\,, \end{aligned} \tag{23}$$

where i, j, k correspond to the diquarks with the spin-parity $J^P = 0^+$, 1^+.

The other choices of point s_0 do not change essentially the contributions of α_l, $l = 1-3$, therefore we omit the indices s_0^{ik}.

We calculate the realistic case of ${}^3He$, containing the u and d quarks. The solutions of the system of 15 equations are considered as $\alpha_i(s) = F_i(s,\lambda_i)/D(s)$, where zeros of $D(s)$ determinant define the mass of bound states (${}^3He$ and ${}^3H$).

The mass of ${}^3He$ is equal to $M = 2809\,MeV$. We propose the masses of p and n states are equal to $M = 940\,MeV$ (without electromagnetic interactions), then the ${}^3H$ mass is equal to the mass of ${}^3He$.

The reduced amplitudes of ${}^3He$ are given in the Appendix A.

IV. CALCULATION RESULTS.

The poles of the reduced amplitudes α_l ($l = 1-3$) correspond to the bound state and determines the mass of the nine-quark state with $J^P = \frac{1}{2}^+$ and isospin $I = \frac{1}{2}$. The quark u, d masses are equal to $m_{u,d} = 410\,MeV$, coincide with the ordinary baryon ones in our model (Sec. II). The model in question has only two parameters: the cutoff parameter $\Lambda = 11$ (similar to the three quark model (Sec. II)) and the gluon coupling constant $g_V = 0.1536$. This parameter is determined by the ${}^3He$ mass $M = 2809\,MeV$. The ${}^3H$ mass is equal to one. In the nuclei region the gluon interaction is smaller in $2-3$ time as compared of the gluon interaction of low-lying hadrons $(0.35-0.45)$.

The estimation of theoretical error on the ${}^3He$ is equal to $1\,MeV$. This results was obtained by the choice of model parameters. We predict the mix pn + 3 quarks, pp + 3 quarks systems and two diquark + 5 quarks.

The functions I_1, I_2, I_3, I_4, I_5, I_6, I_7, I_8, I_9, I_{10} are taken in paper [19]. The other functions are determined by the following formulae (see also Fig. 3):

$$I_{11}(12,34,15,36,47) = I_1(12,15) \times I_2(34,36,47)\,, \tag{24}$$

$$I_{12}(12,34,56,17) = I_1(12,17)\,, \tag{25}$$

$$I_{13}(12,34,56,17,28) = I_2(12,17,28)\,, \tag{26}$$

$$I_{14}(12,34,56,17,38) = I_1(12,17) \times I_1(34,38)\,, \tag{27}$$

$$I_{15}(12,34,56,23,47) = I_7(12,34,23,47)\,, \tag{28}$$

$$I_{16}(12,34,56,17,23,48) = I_8(12,34,17,23,48)\,, \tag{29}$$

$$I_{17}(12,34,56,17,45) = I_1(12,17)\times I_3(34,56,45)\,, \tag{30}$$

$$I_{18}(12,34,56,17,38,49) = I_1(12,17)\times I_2(34,38,49)\,, \tag{31}$$

$$I_{19}(12,34,56,17,38,59) = I_1(12,17)\times I_1(34,38)\times I_1(56,59)\,, \tag{32}$$

$$I_{20}(12,34,56,17,45,68) = I_1(12,17)\times I_7(34,56,45,68)\,, \tag{33}$$

$$I_{21}(12,34,56,17,28,45) = I_2(12,17,28)\times I_3(34,56,45)\,. \tag{34}$$

For the new function I_{22} we used the following equation:

$$\begin{aligned}
I_{22}(ijklmn) \;=\; & \frac{B_l(s_0^{23})B_m(s_0^{45})B_n(s_0^{67})}{B_i(s_0^{12})B_j(s_0^{34})B_k(s_0^{56})} \int\limits_{4m^2}^{\Lambda} \frac{ds'_{12}}{\pi}\frac{G_i^2(s_0^{12})\rho_i(s'_{12})}{s'_{12}-s_0^{12}} \int\limits_{4m^2}^{\Lambda} \frac{ds'_{34}}{\pi}\frac{G_j^2(s_0^{34})\rho_j(s'_{34})}{s'_{34}-s_0^{34}} \\
\times & \int\limits_{4m^2}^{\Lambda} \frac{ds'_{56}}{\pi}\frac{G_k^2(s_0^{56})\rho_k(s'_{56})}{s'_{56}-s_0^{56}} \frac{1}{(2\pi)^2} \int\limits_{-1}^{+1}\frac{dz_1}{2} \int\limits_{-1}^{+1}\frac{dz_2}{2} \int\limits_{-1}^{+1}\frac{dz_3}{2} \int\limits_{-1}^{+1}\frac{dz_4}{2} \int\limits_{-1}^{+1}\frac{dz_6}{2} \int\limits_{-1}^{+1}\frac{dz_7}{2} \\
\times & \int\limits_{z_5^-}^{z_5^+} dz_5 \int\limits_{z_8^-}^{z_8^+} dz_8 \frac{1}{\sqrt{1-z_3^2-z_4^2-z_5^2+2z_3z_4z_5}} \frac{1}{\sqrt{1-z_2^2-z_7^2-z_8^2+2z_2z_7z_8}} \\
\times & \frac{1}{1-B_l(s'_{23})}\frac{1}{1-B_m(s'_{45})}\frac{1}{1-B_n(s'_{67})}\,.
\end{aligned} \tag{35}$$

The contribution of this new functions is about 10^{-8} and the contribution of I_{10}, I_{17}, I_{19}, I_{20}, I_{21} (Eqs. (30), (32) – (34)) are similar to I_{22}. We do not take into account these functions in Appendix A. The $ijklmn$ determine the diquarks contributions.

V. CONCLUSIONS.

The structure and interactions of the light nuclei have been the focus of experimental and theoretical exploration since the infancy of nuclear physics.

The known way with which to calculate the low-energy properties of hadronic and nuclear systems rigorously is Lattice QCD (LQCD). In LQCD calculations, the quark and gluon fields are defined on a discretized space-time of finite volume of the lattice, such deviations can be systematically removed by reducing the lattice spacing, increasing the lattice volume and extrapolating to the continuum and infinite volume limits using the known dependences determined with effective field theory (EFT). Calculation of important quantifies in nuclear physics using LQCD is only now becoming practical, with first calculations of simple multi-baryon interactions being recently performed, although not at the physical values of the light-quark masses. Early exploratory quenched calculations of the NN scattering lengths [35, 36] performed more than a decade ago have been used by $n_f = 2+1$ calculations within the last few years [37, 38] (and added to by futher quenched calculations [39, 40]).

Further, the first quenched calculations of the deuteron [35], ^3He and ^4He [36] have been performed with $n_f = 2+1$ calculations of ^3He and multi-baryon systems containing strange quarks [37]. In addition efforts to explore nuclei and nuclear matter using the strong coupling limit of QCD have led to same interesting observations [38]. Future calculation at smaller spacing and at lighter quark masses will facilitate such extrapolations and lead to first predictions for the spectrum of light nuclei, with completely quantified uncertainties, that can be compared with experiment.

We have considered the dibaryons with the u, d, s-quarks. The H-particle, $N\Omega$-state and di-Ω may be strong interaction stable. Up to now, these three interesting candidates of dibaryons are still not found or confirmed by experiments. It seems that one should go beyond these candidates and should search the possible candidates in a wider region, especially the systems with multistrangeness in terms of a more reliable model.

In our paper the dynamics of quark interactions is defined by the Chew-Mandelstam functions (Table II). We include only two parameters: the cutoff Λ, gluon coupling constant g_V.

The comparison of baryons with the ^3He and ^3H state gluon coupling constants gives rise to $2-3$ time smaller the baryon one.

The quark-gluon interaction allows us to calculate the mass spectrum at light hypernuclei.

Acknowledgments

The authors would like to thank to T. Barnes, L.V. Krasnov for useful discussion.

---

[1] K. Grotz, H.V. Klapdor, and J. Metzinger, Phys. Rev. C**33**, 1263 (1986).
[2] J.J. Arvieux et al., Nucl. Phys. A**431**, 613 (1984).
[3] B. Keister, Phys. Rev. C**24**, 2628 (1981).
[4] M. Rekalo, G. Gakh, and A. Rekalo, Phys. Atom. Nucl. **57**, 698 (1994).
[5] M. Rekalo, G. Gakh, and A. Rekalo, Phys. Atom. Nucl. **60**, 1228 (1994).
[6] C. Perdrisat et al., Phys. Rev. Lett. **59**, 2640 (1987).
[7] M. Dolidze and G. Lykasov, Z. Phys. A**335**, 95 (1990).
[8] L. Azhgirei et al., Phys. Lett. B**361**, 21 (1995).
[9] L. Azhgirei et al., 14th International IUPAP Conference on Few Body Problems in Physics (ICFBP 14).
[10] L. Azhgirei et al., Phys. Lett. B**391**, 22 (1997).
[11] V. Punjabi et al., Phys. Lett. B**350**, 178 (1995).
[12] A. Kerman and L. Kisslinger, Phys. Rev. **180**, 1483 (1969).
[13] A. Illarionov and G. Lykasov, Phys. Rev. C**64**, 044004 (2001).
[14] A. Ierusalimov, G. Lykasov, and M. Viviani, In 20th European Conference on Few-Body Problems in Physics (EFBR 20), Pisa, Italy, 10 – 14 September 2007; Few-Body v.44 pp. 315 – 318 (2008).
[15] A.V. Anisovich and V.V. Anisovich, Yad. Fiz. **53**, 1485 (1991) [Sov. J. Nucl. Phys. **53**, 915 (1991)].
[16] S.M. Gerasyuta, Yad. Fiz. **55**, 3030 (1992) [Sov. J. Nucl. Phys. **55**, 1693 (1992)].
[17] S.M. Gerasyuta, Nuovo Cimento Soc. Ital. Fis. A**106**, 37 (1993).
[18] S.M. Gerasyuta, Z. Phys. C**60**, 683 (1993).
[19] S.M. Gerasyuta and E.E. Matskevich, Phys. Rev. D**82**, 056002 (2010).
[20] G. Chew, S. Mandelstam, Phys. Rev. **119**, 467 (1960).
[21] Y. Nambu and G. Jona-Lasinio, Phys. Rev. **122**, 345 (1961); **124**, 246 (1961).
[22] T. Appelqvist and J.D. Bjorken, Phys. Rev. D**4**, 3726 (1971).
[23] C.C. Chiang, C.B. Chiu, E.C.G. Sudarshan and X. Tata, Phys. Rev. D**25**, 1136 (1982).
[24] A. De Rujula, H. Georgi and S.L. Glashow, Phys. Rev. D**12**, 147 (1975).
[25] V.V. Anisovich, S.M. Gerasyuta and A.V. Sarantsev, Int. J. Mod. Phys. A**6**, 625 (1991).
[26] V.V. Anisovich and A.A. Anselm, Usp. Fiz. Nauk **88**, 287 (1966).
[27] J. Berlinger et al. (Particle Data Group), Phys. Rev. D**86**, 010001 (2012).
[28] G.'t Hooft, Nucl. Phys. B**72**, 461 (1974).
[29] G. Veneziano, Nucl. Phys. B**117**, 519 (1976).
[30] E. Witten, Nucl. Phys. B**160**, 57 (1979).
[31] O.A. Yakubovsky, Sov. J. Nucl. Phys. **5**, 1312 (1967).
[32] S.P. Merkuriev and L.D. Faddeev, Quantum Scattering Theory for System of Few Particles (Nauka, Moscow, 1985) p. 398.
[33] C. Samanta, P. Roy Chowdhury, and D.N. Basu, arXiv: nucl-th/0504085 (2006).
[34] C. Samanta, arXiv: 1003.4227 [nucl-th] (2010).

[35] M. Fukugita, Y. Kuramashi, H. Mino, M. Okawa, and A. Ukawa, Phys. Rev. Lett. **73**, 2176 (1994).
[36] M. Fukugita, Y. Kuramashi, H. Mino, M. Okawa, and A. Ukawa, Phys. Rev. D**52**, 3003 (1996).
[37] S.R. Beane, P.F. Bedaque, K. Orginos, and J.M. Savage, Phys. Rev. Lett. **97**, 012001 (2006).
[38] N. Ishii, S. Aoki, and T. Hatsuda, Phys. Rev. Lett. **99**, 022001 (2007).
[39] S.R. Beane et al., Phys. Rev. D**80**, 074501 (2009).
[40] S.R. Beane et al., arXiv: 1206.5219 [hep-lat] (2012).



Fig. 1. Graphic representation of the equation for the amplitude $A_1(s, s_{12})$.


1^{uu}
1 u
2 u
3 u
4 u
5 u
6 d
7 d
8 d
9 d
=
$\alpha_1^{1^{uu}}$
λ
+ 6
$\alpha_1^{1^{uu}} I_1(1^{uu}1^{uu})$
+ 8
0^{ud}
$\alpha_1^{0^{ud}} I_1(1^{uu}0^{ud})$
+ 6
$\alpha_2^{1^{uu}1^{uu}} I_2(1^{uu}1^{uu}1^{uu})$
+ 24
$\alpha_2^{1^{uu}0^{ud}} I_2(1^{uu}1^{uu}0^{ud})$
+ 12
$\alpha_2^{0^{ud}0^{ud}} I_2(1^{uu}0^{ud}0^{ud})$

Fig. 2. The reduced amplitude $\alpha_1^{1^{uu}}$ with the functions I_i.



α_1

:



λ

;



$\alpha_1\, I_1(12,13)$

;



$\alpha_2\, I_2(12,13,24)$



α_2

:



λ

;



$\alpha_1\, I_3(12,34,23)$

;



$\alpha_1\, I_4(12,34,15)$
$= \alpha_1\, I_1(12,15)$

;



$\alpha_2\, I_5(12,34,15,26)$
$= \alpha_2\, I_2(12,15,26)$

;



$\alpha_2\, I_6(12,34,15,36)$
$= \alpha_2\, I_1(12,15) \times I_1(34,36)$

;



$\alpha_2\, I_7(12,34,23,45)$

;



$\alpha_3\, I_8(12,34,15,23,46)$

;



$\alpha_3\, I_{11}(12,34,15,36,47)$
$= \alpha_3\, I_1(12,15) \times I_2(34,36,47)$



α_3

:



λ

;



$\alpha_1\, I_9(12,34,56,23)$
$= \alpha_1\, I_3(12,34,23)$

;



$\alpha_2\, I_{10}(12,34,56,23,45)$

;



$\alpha_1\, I_{12}(12,34,56,17)$
$= \alpha_1\, I_1(12,17)$

;



$\alpha_2\, I_{13}(12,34,56,17,28)$
$= \alpha_2\, I_2(12,17,28)$

;



$\alpha_2\, I_{14}(12,34,56,17,38)$
$= \alpha_2\, I_1(12,17) \times I_1(34,38)$

;



$\alpha_2\, I_{15}(12,34,56,17,23,47)$
$= \alpha_2\, I_7(12,34,17,23,47)$

;



$\alpha_3\, I_{16}(12,34,56,17,23,48)$
$= \alpha_3\, I_8(12,34,17,23,48)$

;



$\alpha_2\, I_{17}(12,34,56,17,45)$
$= \alpha_2\, I_1(12,17) \times I_3(34,56,45)$

;



$\alpha_3\, I_{18}(12,34,56,17,38,49)$
$= \alpha_3\, I_1(12,17) \times I_2(34,38,49)$

;



$\alpha_3\, I_{19}(12,34,56,17,38,59)$
$= \alpha_3\, I_1(12,17) \times I_1(34,38)$
$\times I_1(56,59)$

;



$\alpha_3\, I_{20}(12,34,56,17,45,68)$
$= \alpha_3\, I_1(12,17) \times I_7(34,56,45,68)$

;



$\alpha_3\, I_{21}(12,34,56,17,28,45)$
$= \alpha_3\, I_2(12,17,28) \times I_3(34,56,45)$

$\alpha_3\, I_{22}(12,34,56,23,45,67)$

Fig. 3. Contributions of different diagrams to the total amplitudes.

Appendix A

The coupled equations for the reduced amplitudes of ^3He:

$$\begin{aligned}
\alpha_1^{1^{uu}} &= \lambda + 6\,\alpha_1^{1^{uu}} I_1(1^{uu}1^{uu}) + 8\,\alpha_1^{0^{ud}} I_1(1^{uu}0^{ud}) + 6\,\alpha_2^{1^{uu}1^{uu}} I_2(1^{uu}1^{uu}1^{uu}) + 24\,\alpha_2^{1^{uu}0^{ud}} I_2(1^{uu}1^{uu}0^{ud}) \\
&+ 12\,\alpha_2^{0^{ud}0^{ud}} I_2(1^{uu}0^{ud}0^{ud})
\end{aligned} \tag{A1}$$

$$\begin{aligned}
\alpha_1^{0^{ud}} &= \lambda + 4\,\alpha_1^{1^{uu}} I_1(0^{ud}1^{uu}) + 7\,\alpha_1^{0^{ud}} I_1(0^{ud}0^{ud}) + 3\,\alpha_1^{1^{dd}} I_1(0^{ud}1^{dd}) + 12\,\alpha_2^{1^{uu}0^{ud}} I_2(0^{ud}1^{uu}0^{ud}) \\
&+ 12\,\alpha_2^{1^{uu}1^{dd}} I_2(0^{ud}1^{uu}1^{dd}) + 12\,\alpha_2^{0^{ud}0^{ud}} I_2(0^{ud}0^{ud}0^{ud}) + 6\,\alpha_2^{1^{dd}0^{ud}} I_2(0^{ud}0^{ud}1^{dd})
\end{aligned} \tag{A2}$$

$$\begin{aligned}
\alpha_1^{1^{dd}} &= \lambda + 10\,\alpha_1^{0^{ud}} I_1(1^{dd}0^{ud}) + 4\,\alpha_1^{1^{dd}} I_1(1^{dd}1^{dd}) + 20\,\alpha_2^{0^{ud}0^{ud}} I_2(1^{dd}0^{ud}0^{ud}) + 20\,\alpha_2^{1^{dd}0^{ud}} I_2(1^{dd}1^{dd}0^{ud}) \\
&+ 2\,\alpha_2^{1^{dd}1^{dd}} I_2(1^{dd}1^{dd}1^{dd})
\end{aligned} \tag{A3}$$

$$\begin{aligned}
\alpha_2^{1^{uu}1^{uu}} &= \lambda + \alpha_1^{1^{uu}} (4\,I_3(1^{uu}1^{uu}1^{uu}) + 4\,I_4(1^{uu}1^{uu}1^{uu})) + 16\,\alpha_1^{0^{ud}} I_4(1^{uu}1^{uu}0^{ud}) \\
&+ 8\,\alpha_2^{1^{uu}1^{uu}} I_7(1^{uu}1^{uu}1^{uu}1^{uu}) + \alpha_2^{1^{uu}0^{ud}} (16\,I_5(1^{uu}1^{uu}1^{uu}0^{ud}) + 32\,I_6(1^{uu}1^{uu}1^{uu}0^{ud}) \\
&+ 32\,I_7(1^{uu}1^{uu}1^{uu}0^{ud})) + \alpha_2^{0^{ud}0^{ud}} (24\,I_5(1^{uu}1^{uu}0^{ud}0^{ud}) + 48\,I_6(1^{uu}1^{uu}0^{ud}0^{ud})) \\
&+ 32\,\alpha_3^{1^{uu}1^{uu}0^{ud}} I_8(1^{uu}1^{uu}1^{uu}1^{uu}0^{ud}) + \alpha_3^{1^{uu}0^{ud}0^{ud}} (48\,I_8(1^{uu}1^{uu}0^{ud}1^{uu}0^{ud}) \\
&+ 48\,I_{11}(1^{uu}1^{uu}1^{uu}0^{ud}0^{ud}) + 96\,I_{11}(1^{uu}1^{uu}0^{ud}1^{uu}0^{ud})) + 96\,\alpha_3^{0^{ud}0^{ud}0^{ud}} I_{11}(1^{uu}1^{uu}0^{ud}0^{ud}0^{ud})
\end{aligned} \tag{A4}$$

$$\begin{aligned}
\alpha_2^{1^{uu}0^{ud}} &= \lambda + \alpha_1^{1^{uu}} (2\,I_3(1^{uu}0^{ud}1^{uu}) + 4\,I_4(1^{uu}0^{ud}1^{uu}) + 2\,I_4(0^{ud}1^{uu}1^{uu})) + \alpha_1^{0^{ud}} (2\,I_3(1^{uu}0^{ud}0^{ud}) \\
&+ 6\,I_4(1^{uu}0^{ud}0^{ud}) + 5\,I_4(0^{ud}1^{uu}0^{ud})) + 3\,\alpha_1^{1^{dd}} I_4(0^{ud}1^{uu}1^{dd}) + \alpha_2^{1^{uu}1^{uu}} (2\,I_5(1^{uu}0^{ud}1^{uu}1^{uu}) \\
&+ 4\,I_6(1^{uu}0^{ud}1^{uu}1^{uu}) + 4\,I_7(0^{ud}1^{uu}1^{uu}1^{uu})) + \alpha_2^{1^{uu}0^{ud}} (12\,I_5(1^{uu}0^{ud}1^{uu}0^{ud}) + 2\,I_5(0^{ud}1^{uu}1^{uu}0^{ud}) \\
&+ 16\,I_6(1^{uu}0^{ud}1^{uu}0^{ud}) + 12\,I_6(1^{uu}0^{ud}0^{ud}1^{uu}) + 4\,I_7(1^{uu}0^{ud}1^{uu}0^{ud}) + 4\,I_7(1^{uu}0^{ud}0^{ud}1^{uu}) \\
&+ 6\,I_7(0^{ud}1^{uu}1^{uu}0^{ud}) + 4\,I_7(0^{ud}1^{uu}0^{ud}1^{uu})) + \alpha_2^{0^{ud}0^{ud}} (6\,I_5(1^{uu}0^{ud}0^{ud}0^{ud}) + 6\,I_5(0^{ud}1^{uu}0^{ud}0^{ud}) \\
&+ 24\,I_6(1^{uu}0^{ud}0^{ud}0^{ud}) + 6\,I_7(1^{uu}0^{ud}0^{ud}0^{ud}) + 6\,I_7(0^{ud}1^{uu}0^{ud}0^{ud})) + \alpha_2^{1^{uu}1^{dd}} (6\,I_5(0^{ud}1^{uu}1^{uu}1^{dd}) \\
&+ 12\,I_6(1^{uu}0^{ud}1^{uu}1^{dd}) + 6\,I_7(1^{uu}0^{ud}1^{uu}1^{dd})) + \alpha_2^{1^{dd}0^{ud}} (6\,I_5(0^{ud}1^{uu}0^{ud}1^{dd}) + 12\,I_6(1^{uu}0^{ud}0^{ud}1^{dd})) \\
&+ \alpha_3^{1^{uu}1^{uu}0^{ud}} (4\,I_8(1^{uu}0^{ud}1^{uu}1^{uu}0^{ud}) + 4\,I_8(1^{uu}0^{ud}1^{uu}0^{ud}1^{uu}) + 6\,I_{11}(0^{ud}1^{uu}0^{ud}1^{uu}1^{uu}) \\
&+ 12\,I_{11}(0^{ud}1^{uu}1^{uu}1^{uu}0^{ud})) + \alpha_3^{1^{uu}1^{uu}1^{dd}} (12\,I_8(1^{uu}0^{ud}1^{uu}1^{uu}1^{dd}) + 12\,I_{11}(1^{uu}0^{ud}1^{uu}1^{uu}1^{dd}) \\
&+ 6\,I_{11}(0^{ud}1^{uu}1^{dd}1^{uu}1^{uu}) + \alpha_3^{1^{uu}0^{ud}0^{ud}} (12\,I_8(1^{uu}0^{ud}0^{ud}1^{uu}0^{ud}) + 12\,I_8(1^{uu}0^{ud}1^{uu}0^{ud}0^{ud}) \\
&+ 12\,I_8(1^{uu}0^{ud}0^{ud}0^{ud}1^{uu}) + 12\,I_{11}(1^{uu}0^{ud}1^{uu}0^{ud}0^{ud}) + 12\,I_{11}(1^{uu}0^{ud}0^{ud}1^{uu}0^{ud}) \\
&+ 12\,I_{11}(0^{ud}1^{uu}1^{uu}0^{ud}0^{ud}) + 36\,I_{11}(0^{ud}1^{uu}0^{ud}1^{uu}0^{ud})) + \alpha_3^{1^{uu}0^{ud}1^{dd}} (12\,I_8(1^{uu}0^{ud}0^{ud}1^{uu}1^{dd}) \\
&+ 24\,I_{11}(1^{uu}0^{ud}0^{ud}1^{uu}1^{dd}) + 24\,I_{11}(1^{uu}0^{ud}1^{uu}0^{ud}1^{dd}) + 24\,I_{11}(0^{ud}1^{uu}1^{dd}1^{uu}0^{ud})) \\
&+ \alpha_3^{0^{ud}0^{ud}0^{ud}} (12\,I_8(1^{uu}0^{ud}0^{ud}0^{ud}0^{ud}) + 24\,I_{11}(1^{uu}0^{ud}0^{ud}0^{ud}0^{ud}) + 18\,I_{11}(0^{ud}1^{uu}0^{ud}0^{ud}0^{ud})) \\
&+ \alpha_3^{0^{ud}0^{ud}1^{dd}} (12\,I_{11}(1^{uu}0^{ud}0^{ud}0^{ud}1^{dd}) + 6\,I_{11}(0^{ud}1^{uu}1^{dd}0^{ud}0^{ud}))
\end{aligned} \tag{A5}$$

$$\begin{aligned}
\alpha_2^{0^{ud}0^{ud}} &= \lambda + \alpha_1^{1^{uu}} (I_3(0^{ud}0^{ud}1^{uu}) + 6\,I_4(0^{ud}0^{ud}1^{uu})) + \alpha_1^{0^{ud}} (2\,I_3(0^{ud}0^{ud}0^{ud}) + 10\,I_4(0^{ud}0^{ud}0^{ud})) \\
&+ \alpha_1^{1^{dd}} (I_3(0^{ud}0^{ud}1^{dd}) + 4\,I_4(0^{ud}0^{ud}1^{dd})) + 6\,\alpha_2^{1^{uu}1^{uu}} I_6(0^{ud}0^{ud}1^{uu}1^{uu}) \\
&+ \alpha_2^{1^{uu}0^{ud}} (12\,I_5(0^{ud}0^{ud}1^{uu}0^{ud}) + 24\,I_6(0^{ud}0^{ud}1^{uu}0^{ud}) + 6\,I_7(0^{ud}0^{ud}1^{uu}0^{ud}) + 6\,I_7(0^{ud}0^{ud}0^{ud}1^{uu})) \\
&+ \alpha_2^{0^{ud}0^{ud}} (12\,I_5(0^{ud}0^{ud}0^{ud}0^{ud}) + 20\,I_6(0^{ud}0^{ud}0^{ud}0^{ud}) + 10\,I_7(0^{ud}0^{ud}0^{ud}0^{ud})) \\
&+ \alpha_2^{1^{uu}1^{dd}} (12\,I_5(0^{ud}0^{ud}1^{uu}1^{dd}) + 12\,I_6(0^{ud}0^{ud}1^{uu}1^{dd}) + 6\,I_7(0^{ud}0^{ud}1^{dd}1^{uu}) + 4\,I_7(0^{ud}0^{ud}1^{uu}1^{dd})) \\
&+ \alpha_2^{1^{dd}0^{ud}} (4\,I_5(0^{ud}0^{ud}0^{ud}1^{dd}) + 16\,I_6(0^{ud}0^{ud}0^{ud}1^{dd}) + 4\,I_7(0^{ud}0^{ud}0^{ud}1^{dd}) + 4\,I_7(0^{ud}0^{ud}1^{dd}0^{ud})) \\
&+ 2\,\alpha_2^{1^{dd}1^{dd}} I_6(0^{ud}0^{ud}1^{dd}1^{dd}) + 12\,\alpha_3^{1^{uu}1^{uu}0^{ud}} I_{11}(0^{ud}0^{ud}1^{uu}1^{uu}0^{ud}) + \alpha_3^{1^{uu}1^{uu}1^{dd}} (6\,I_8(0^{ud}0^{ud}1^{uu}1^{dd}1^{uu}) \\
&+ 24\,I_{11}(0^{ud}0^{ud}1^{uu}1^{uu}1^{dd})) + \alpha_3^{1^{uu}0^{ud}0^{ud}} (12\,I_8(0^{ud}0^{ud}1^{uu}0^{ud}0^{ud}) + 6\,I_8(0^{ud}0^{ud}0^{ud}1^{uu}0^{ud}) \\
&+ 24\,I_{11}(0^{ud}0^{ud}1^{uu}0^{ud}0^{ud}) + 36\,I_{11}(0^{ud}0^{ud}0^{ud}1^{uu}0^{ud})) + \alpha_3^{1^{uu}0^{ud}1^{dd}} (12\,I_8(0^{ud}0^{ud}1^{uu}0^{ud}1^{dd}) \\
&+ 12\,I_8(0^{ud}0^{ud}1^{uu}1^{dd}0^{ud}) + 12\,I_8(0^{ud}0^{ud}0^{ud}1^{uu}1^{dd}) + 12\,I_{11}(0^{ud}0^{ud}1^{uu}0^{ud}1^{dd}) \\
&+ 36\,I_{11}(0^{ud}0^{ud}0^{ud}1^{uu}1^{dd}) + 24\,I_{11}(0^{ud}0^{ud}1^{dd}1^{uu}0^{ud})) + \alpha_3^{0^{ud}0^{ud}0^{ud}} (12\,I_8(0^{ud}0^{ud}0^{ud}0^{ud}0^{ud})
\end{aligned}$$

$$
\begin{aligned}
&+\ 36\, I_{11}(0^{ud}0^{ud}0^{ud}0^{ud}0^{ud})) + \alpha_3^{0^{ud}0^{ud}1^{dd}}(4\, I_8(0^{ud}0^{ud}0^{ud}0^{ud}1^{dd}) + 2\, I_8(0^{ud}0^{ud}0^{ud}1^{dd}0^{ud}) \\
&+\ 12\, I_{11}(0^{ud}0^{ud}0^{ud}0^{ud}1^{dd}) + 12\, I_{11}(0^{ud}0^{ud}1^{dd}0^{ud}0^{ud}))
\end{aligned} \tag{A6}
$$

$$
\begin{aligned}
\alpha_2^{1^{uu}1^{dd}} &= \lambda + 6\, \alpha_1^{1^{uu}} I_4(1^{uu}1^{dd}1^{uu}) + \alpha_1^{0^{ud}}(4\, I_3(1^{uu}1^{dd}0^{ud}) + 4\, I_4(1^{uu}1^{dd}0^{ud}) + 6\, I_4(1^{dd}1^{uu}0^{ud})) \\
&+\ 4\, \alpha_1^{1^{dd}} I_4(1^{dd}1^{uu}1^{dd}) + 6\, \alpha_2^{1^{uu}1^{uu}} I_5(1^{uu}1^{dd}1^{uu}1^{uu}) + \alpha_2^{1^{uu}0^{ud}}(12\, I_5(1^{uu}1^{dd}1^{uu}0^{ud}) \\
&+\ 24\, I_6(1^{uu}1^{dd}1^{uu}0^{ud}) + 12\, I_7(1^{dd}1^{uu}0^{ud}1^{uu})) + \alpha_2^{0^{ud}0^{ud}}(2\, I_5(1^{uu}1^{dd}0^{ud}0^{ud}) + 6\, I_5(1^{dd}1^{uu}0^{ud}0^{ud}) \\
&+\ 24\, I_6(1^{uu}1^{dd}0^{ud}0^{ud}) + 12\, I_7(1^{uu}1^{dd}0^{ud}0^{ud}) + 8\, I_7(1^{dd}1^{uu}0^{ud}0^{ud})) + 24\, \alpha_2^{1^{uu}1^{dd}} I_6(1^{uu}1^{dd}1^{uu}1^{dd}) \\
&+\ \alpha_2^{1^{dd}0^{ud}}(12\, I_5(1^{dd}1^{uu}0^{ud}1^{dd}) + 8\, I_6(1^{uu}1^{dd}0^{ud}1^{dd}) + 8\, I_7(1^{uu}1^{dd}0^{ud}1^{dd})) + 2\, \alpha_2^{1^{dd}1^{dd}} I_5(1^{dd}1^{uu}1^{dd}1^{dd}) \\
&+\ 12\, \alpha_3^{1^{uu}1^{uu}0^{ud}} I_{11}(1^{dd}1^{uu}0^{ud}1^{uu}1^{uu}) + 24\, \alpha_3^{1^{uu}1^{uu}1^{dd}} I_{11}(1^{dd}1^{uu}1^{dd}1^{uu}1^{uu}) \\
&+\ \alpha_3^{1^{uu}0^{ud}0^{ud}}(24\, I_8(1^{uu}1^{dd}1^{uu}0^{ud}0^{ud}) + 12\, I_{11}(1^{uu}1^{dd}1^{uu}0^{ud}0^{ud}) + 48\, I_{11}(1^{dd}1^{uu}0^{ud}1^{uu}0^{ud})) \\
&+\ \alpha_3^{1^{uu}0^{ud}1^{dd}}(24\, I_8(1^{uu}1^{dd}1^{uu}0^{ud}1^{dd}) + 48\, I_{11}(1^{uu}1^{dd}1^{uu}0^{ud}1^{dd}) + 24\, I_{11}(1^{dd}1^{uu}1^{dd}1^{uu}0^{ud})) \\
&+\ \alpha_3^{0^{ud}0^{ud}0^{ud}}(24\, I_8(1^{uu}1^{dd}0^{ud}0^{ud}0^{ud}) + 24\, I_{11}(1^{uu}1^{dd}0^{ud}0^{ud}0^{ud}) + 12\, I_{11}(1^{dd}1^{uu}0^{ud}0^{ud}0^{ud})) \\
&+\ \alpha_3^{0^{ud}0^{ud}1^{dd}}(8\, I_8(1^{uu}1^{dd}0^{ud}0^{ud}1^{dd}) + 24\, I_{11}(1^{uu}1^{dd}0^{ud}0^{ud}1^{dd}))
\end{aligned} \tag{A7}
$$

$$
\begin{aligned}
\alpha_2^{1^{dd}0^{ud}} &= \lambda + 4\, \alpha_1^{1^{uu}} I_4(0^{ud}1^{dd}1^{uu}) + \alpha_1^{0^{ud}}(2\, I_3(1^{dd}0^{ud}0^{ud}) + 8\, I_4(1^{dd}0^{ud}0^{ud}) + 5\, I_4(0^{ud}1^{dd}0^{ud})) \\
&+\ \alpha_1^{1^{dd}}(2\, I_3(1^{dd}0^{ud}1^{dd}) + 2\, I_4(1^{dd}0^{ud}1^{dd}) + I_4(0^{ud}1^{dd}1^{dd})) + \alpha_2^{1^{uu}0^{ud}}(12\, I_5(0^{ud}1^{dd}1^{uu}0^{ud}) \\
&+\ 24\, I_6(1^{dd}0^{ud}0^{ud}1^{uu})) + \alpha_2^{0^{ud}0^{ud}}(12\, I_5(1^{dd}0^{ud}0^{ud}0^{ud}) + 4\, I_5(0^{ud}1^{dd}0^{ud}0^{ud}) + 32\, I_6(1^{dd}0^{ud}0^{ud}0^{ud}) \\
&+\ 8\, I_7(1^{dd}0^{ud}0^{ud}0^{ud}) + 8\, I_7(0^{ud}1^{dd}0^{ud}0^{ud})) + \alpha_2^{1^{uu}1^{dd}}(4\, I_5(0^{ud}1^{dd}1^{uu}1^{dd}) + 8\, I_6(1^{dd}0^{ud}1^{dd}1^{uu}) \\
&+\ 8\, I_7(1^{dd}0^{ud}1^{dd}1^{uu})) + \alpha_2^{1^{dd}0^{ud}}(8\, I_5(1^{dd}0^{ud}0^{ud}1^{dd}) + 8\, I_6(1^{dd}0^{ud}0^{ud}1^{dd}) + 8\, I_6(1^{dd}0^{ud}1^{dd}0^{ud}) \\
&+\ 2\, I_7(1^{dd}0^{ud}0^{ud}1^{dd}) + 2\, I_7(1^{dd}0^{ud}1^{dd}0^{ud}) + 8\, I_7(0^{ud}1^{dd}1^{dd}0^{ud}) + 2\, I_7(0^{ud}1^{dd}0^{ud}1^{dd})) \\
&+\ 2\, \alpha_2^{1^{dd}1^{dd}} I_7(0^{ud}1^{dd}1^{dd}1^{dd}) + \alpha_3^{1^{uu}0^{ud}0^{ud}}(48\, I_{11}(1^{dd}0^{ud}0^{ud}1^{uu}0^{ud}) + 24\, I_{11}(0^{ud}1^{dd}1^{uu}0^{ud}0^{ud})) \\
&+\ \alpha_3^{1^{uu}0^{ud}1^{dd}}(24\, I_8(1^{dd}0^{ud}0^{ud}1^{dd}1^{uu}) + 24\, I_{11}(1^{dd}0^{ud}1^{dd}1^{uu}0^{ud}) + 24\, I_{11}(1^{dd}0^{ud}0^{ud}1^{uu}1^{dd}) \\
&+\ 24\, I_{11}(0^{ud}1^{dd}1^{uu}0^{ud}1^{dd})) + \alpha_3^{0^{ud}0^{ud}0^{ud}}(24\, I_8(1^{dd}0^{ud}0^{ud}0^{ud}0^{ud}) + 24\, I_{11}(1^{dd}0^{ud}0^{ud}0^{ud}0^{ud}) \\
&+\ 36\, I_{11}(0^{ud}1^{dd}0^{ud}0^{ud}0^{ud})) + \alpha_3^{0^{ud}0^{ud}1^{dd}}(8\, I_8(1^{dd}0^{ud}0^{ud}0^{ud}1^{dd}) + 8\, I_8(1^{dd}0^{ud}1^{dd}0^{ud}0^{ud}) \\
&+\ 8\, I_8(1^{dd}0^{ud}0^{ud}1^{dd}0^{ud}) + 12\, I_{11}(0^{ud}1^{dd}1^{dd}0^{ud}0^{ud}) + 24\, I_{11}(0^{ud}1^{dd}0^{ud}0^{ud}1^{dd}))
\end{aligned} \tag{A8}
$$

$$
\begin{aligned}
\alpha_2^{1^{dd}1^{dd}} &= \lambda + 20\, \alpha_1^{0^{ud}} I_4(1^{dd}1^{dd}0^{ud}) + 4\, \alpha_1^{1^{dd}} I_3(1^{dd}1^{dd}1^{dd}) + \alpha_2^{0^{ud}0^{ud}}(40\, I_5(1^{dd}1^{dd}0^{ud}0^{ud}) \\
&+\ 80\, I_6(1^{dd}1^{dd}0^{ud}0^{ud})) + 40\, \alpha_2^{1^{dd}0^{ud}} I_7(1^{dd}1^{dd}0^{ud}0^{ud}) + 240\, \alpha_3^{0^{ud}0^{ud}0^{ud}} I_{11}(1^{dd}1^{dd}0^{ud}0^{ud}0^{ud}) \\
&+\ 80\, \alpha_3^{0^{ud}0^{ud}1^{dd}} I_8(1^{dd}1^{dd}0^{ud}1^{dd}0^{ud})
\end{aligned} \tag{A9}
$$

$$
\begin{aligned}
\alpha_3^{1^{uu}1^{uu}0^{ud}} &= \lambda + \alpha_1^{1^{uu}}(4\, I_9(1^{uu}1^{uu}0^{ud}1^{uu}) + 4\, I_9(1^{uu}0^{ud}1^{uu}1^{uu})) + \alpha_1^{0^{ud}}(4\, I_9(1^{uu}0^{ud}1^{uu}0^{ud}) \\
&+\ 12\, I_{12}(1^{uu}1^{uu}0^{ud}0^{ud}) + 3\, I_{12}(0^{ud}1^{uu}1^{uu}0^{ud})) + 3\, \alpha_1^{1^{dd}} I_{12}(0^{ud}1^{uu}1^{uu}1^{dd}) \\
&+\ \alpha_2^{1^{uu}0^{ud}}(24\, I_{15}(1^{uu}1^{uu}0^{ud}1^{uu}0^{ud}) + 12\, I_{15}(0^{ud}1^{uu}1^{uu}1^{uu}0^{ud})) + \alpha_2^{0^{ud}0^{ud}}(12\, I_{13}(1^{uu}1^{uu}0^{ud}0^{ud}0^{ud}) \\
&+\ 24\, I_{14}(1^{uu}1^{uu}0^{ud}0^{ud}0^{ud}) + 24\, I_{14}(1^{uu}0^{ud}1^{uu}0^{ud}0^{ud}) + 12\, I_{15}(1^{uu}0^{ud}1^{uu}0^{ud}0^{ud}) \\
&+\ 12\, I_{15}(0^{ud}1^{uu}1^{uu}0^{ud}0^{ud})) + 12\, \alpha_2^{1^{uu}1^{dd}} I_{15}(1^{uu}0^{ud}1^{uu}1^{uu}1^{dd}) + \alpha_2^{1^{dd}0^{ud}}(6\, I_{13}(0^{ud}1^{uu}1^{uu}0^{ud}1^{dd}) \\
&+\ 24\, I_{14}(1^{uu}0^{ud}1^{uu}0^{ud}1^{dd})) + 24\, \alpha_3^{1^{uu}0^{ud}0^{ud}} I_{16}(1^{uu}1^{uu}0^{ud}0^{ud}1^{uu}0^{ud}) \\
&+\ 24\, \alpha_3^{1^{uu}0^{ud}1^{dd}} I_{16}(1^{uu}0^{ud}1^{uu}0^{ud}1^{uu}1^{dd}) + \alpha_3^{0^{ud}0^{ud}0^{ud}}(24\, I_{16}(1^{uu}0^{ud}1^{uu}0^{ud}0^{ud}0^{ud}) \\
&+\ 12\, I_{18}(0^{ud}1^{uu}1^{uu}0^{ud}0^{ud}0^{ud}) + 24\, I_{18}(1^{uu}1^{uu}0^{ud}0^{ud}0^{ud}0^{ud})) \\
&+\ \alpha_3^{0^{ud}0^{ud}1^{dd}}(12\, I_{18}(0^{ud}1^{uu}1^{uu}1^{dd}0^{ud}0^{ud}) + 24\, I_{18}(1^{uu}0^{ud}1^{uu}0^{ud}0^{ud}1^{dd}))
\end{aligned} \tag{A10}
$$

$$
\begin{aligned}
\alpha_3^{1^{uu}1^{uu}1^{dd}} &= \lambda + 4\, \alpha_1^{1^{uu}} I_9(1^{uu}1^{uu}1^{dd}1^{uu}) + \alpha_1^{0^{ud}}(8\, I_9(1^{uu}1^{dd}1^{uu}0^{ud}) + 8\, I_{12}(1^{uu}1^{uu}1^{dd}0^{ud}) \\
&+\ 2\, I_{12}(1^{dd}1^{uu}1^{uu}0^{ud})) + \alpha_2^{1^{uu}0^{ud}}(8\, I_{13}(1^{uu}1^{uu}1^{dd}1^{uu}0^{ud}) + 16\, I_{15}(1^{uu}1^{uu}1^{dd}1^{uu}0^{ud})) \\
&+\ \alpha_2^{0^{ud}0^{ud}}(4\, I_{13}(1^{uu}1^{uu}1^{dd}0^{ud}0^{ud}) + 8\, I_{14}(1^{uu}1^{uu}1^{dd}0^{ud}0^{ud}) + 16\, I_{14}(1^{uu}1^{dd}1^{uu}0^{ud}0^{ud}) \\
&+\ 8\, I_{15}(1^{uu}1^{dd}1^{uu}0^{ud}0^{ud}) + 16\, I_{15}(1^{dd}1^{uu}1^{uu}0^{ud}0^{ud})) + 4\, \alpha_2^{1^{dd}0^{ud}} I_{13}(1^{dd}1^{uu}1^{uu}0^{ud}1^{dd}) \\
&+\ \alpha_3^{1^{uu}0^{ud}0^{ud}}(8\, I_{16}(1^{uu}1^{uu}1^{dd}0^{ud}1^{uu}0^{ud}) + 16\, I_{18}(1^{uu}1^{uu}1^{dd}0^{ud}1^{uu}0^{ud}))
\end{aligned}
$$

$$\begin{aligned}
&+\ \alpha_3^{0^{ud}0^{ud}0^{ud}}(16\,I_{16}(1^{uu}1^{dd}1^{uu}0^{ud}0^{ud}0^{ud}) + 8\,I_{18}(1^{dd}1^{uu}1^{uu}0^{ud}0^{ud}0^{ud}))\\
&+\ 16\,\alpha_3^{0^{ud}0^{ud}1^{dd}} I_{18}(1^{uu}1^{dd}1^{uu}0^{ud}0^{ud}1^{dd})
\end{aligned}\tag{A11}$$

$$\begin{aligned}
\alpha_3^{1^{uu}0^{ud}0^{ud}} &= \lambda + \alpha_1^{1^{uu}}(I_9(0^{ud}0^{ud}1^{uu}1^{uu}) + 4\,I_9(1^{uu}0^{ud}0^{ud}1^{uu}) + 2\,I_{12}(0^{ud}1^{uu}0^{ud}1^{uu})) + \alpha_1^{0^{ud}}(2\,I_9(0^{ud}0^{ud}1^{uu}0^{ud})\\
&+\ 4\,I_9(1^{uu}0^{ud}0^{ud}0^{ud}) + 4\,I_{12}(1^{uu}0^{ud}0^{ud}0^{ud}) + 6\,I_{12}(0^{ud}1^{uu}0^{ud}0^{ud})) + \alpha_1^{1^{dd}}(I_9(0^{ud}0^{ud}1^{uu}1^{dd})\\
&+\ 4\,I_{12}(0^{ud}1^{uu}0^{ud}1^{dd})) + \alpha_2^{1^{uu}0^{ud}}(4\,I_{13}(1^{uu}0^{ud}0^{ud}1^{uu}0^{ud}) + 4\,I_{14}(0^{ud}0^{ud}1^{uu}1^{uu}0^{ud})\\
&+\ 8\,I_{14}(1^{uu}0^{ud}0^{ud}0^{ud}1^{uu}) + 2\,I_{15}(0^{ud}0^{ud}1^{uu}1^{uu}0^{ud}) + 2\,I_{15}(0^{ud}0^{ud}1^{uu}0^{ud}1^{uu})\\
&+\ 4\,I_{15}(1^{uu}0^{ud}0^{ud}1^{uu}0^{ud}) + 4\,I_{15}(1^{uu}0^{ud}0^{ud}0^{ud}1^{uu}) + 8\,I_{15}(0^{ud}1^{uu}0^{ud}1^{uu}0^{ud}))\\
&+\ \alpha_2^{0^{ud}0^{ud}}(2\,I_{13}(1^{uu}0^{ud}0^{ud}0^{ud}0^{ud}) + 4\,I_{13}(0^{ud}1^{uu}0^{ud}0^{ud}0^{ud}) + 4\,I_{14}(0^{ud}0^{ud}1^{uu}0^{ud}0^{ud})\\
&+\ 16\,I_{14}(1^{uu}0^{ud}0^{ud}0^{ud}0^{ud}) + 6\,I_{15}(0^{ud}0^{ud}1^{uu}0^{ud}0^{ud}) + 8\,I_{15}(1^{uu}0^{ud}0^{ud}0^{ud}0^{ud})\\
&+\ 8\,I_{15}(0^{ud}1^{uu}0^{ud}0^{ud}0^{ud})) + \alpha_2^{1^{uu}1^{dd}}(4\,I_{13}(0^{ud}1^{uu}0^{ud}1^{uu}1^{dd}) + 4\,I_{14}(0^{ud}0^{ud}1^{uu}1^{uu}1^{dd})\\
&+\ 2\,I_{15}(0^{ud}0^{ud}1^{uu}1^{dd}1^{uu}) + 4\,I_{15}(0^{ud}0^{ud}1^{uu}1^{uu}1^{dd}) + 8\,I_{15}(1^{uu}0^{ud}0^{ud}1^{uu}1^{dd}))\\
&+\ \alpha_2^{1^{dd}0^{ud}}(4\,I_{13}(0^{ud}1^{uu}0^{ud}0^{ud}1^{dd}) + 4\,I_{14}(0^{ud}0^{ud}1^{uu}0^{ud}1^{dd}) + 8\,I_{14}(1^{uu}0^{ud}0^{ud}0^{ud}1^{dd})\\
&+\ 4\,I_{15}(0^{ud}0^{ud}1^{uu}0^{ud}1^{dd}) + 4\,I_{15}(0^{ud}0^{ud}1^{uu}1^{dd}0^{ud})) + \alpha_3^{1^{uu}0^{ud}0^{ud}}(8\,I_{16}(1^{uu}0^{ud}0^{ud}0^{ud}0^{ud}1^{uu})\\
&+\ 8\,I_{16}(1^{uu}0^{ud}0^{ud}0^{ud}1^{uu}0^{ud}) + 4\,I_{18}(0^{ud}1^{uu}0^{ud}1^{uu}0^{ud}0^{ud}) + 8\,I_{18}(0^{ud}1^{uu}0^{ud}0^{ud}1^{uu}0^{ud}))\\
&+\ \alpha_3^{1^{uu}0^{ud}1^{dd}}(4\,I_{16}(0^{ud}0^{ud}1^{uu}0^{ud}1^{uu}1^{dd}) + 4\,I_{16}(0^{ud}0^{ud}1^{uu}1^{uu}1^{dd}0^{ud}) + 4\,I_{16}(0^{ud}0^{ud}1^{uu}1^{dd}0^{ud}1^{uu})\\
&+\ 8\,I_{16}(1^{uu}0^{ud}0^{ud}0^{ud}1^{uu}1^{dd}) + 8\,I_{18}(0^{ud}1^{uu}0^{ud}1^{dd}1^{uu}0^{ud}) + 8\,I_{18}(1^{uu}0^{ud}0^{ud}0^{ud}1^{uu}1^{dd})\\
&+\ 4\,I_{18}(0^{ud}0^{ud}1^{uu}0^{ud}1^{uu}1^{dd}) + 4\,I_{18}(0^{ud}0^{ud}1^{uu}1^{uu}0^{ud}1^{dd})) + \alpha_3^{0^{ud}0^{ud}0^{ud}}(4\,I_{16}(0^{ud}0^{ud}1^{uu}0^{ud}0^{ud}0^{ud})\\
&+\ 8\,I_{16}(1^{uu}0^{ud}0^{ud}0^{ud}0^{ud}0^{ud}) + 4\,I_{18}(0^{ud}1^{uu}0^{ud}0^{ud}0^{ud}0^{ud}) + 8\,I_{18}(1^{uu}0^{ud}0^{ud}0^{ud}0^{ud}0^{ud})\\
&+\ 4\,I_{18}(0^{ud}0^{ud}1^{uu}0^{ud}0^{ud}0^{ud})) + \alpha_3^{0^{ud}0^{ud}1^{dd}}(4\,I_{18}(0^{ud}0^{ud}1^{uu}1^{dd}0^{ud}0^{ud})\\
&+\ 4\,I_{18}(0^{ud}0^{ud}1^{uu}0^{ud}0^{ud}1^{dd}))
\end{aligned}\tag{A12}$$

$$\begin{aligned}
\alpha_3^{1^{uu}0^{ud}1^{dd}} &= \lambda + \alpha_1^{1^{uu}}(2\,I_9(1^{uu}0^{ud}1^{dd}1^{uu}) + 2\,I_{12}(0^{ud}1^{uu}1^{dd}1^{uu})) + \alpha_1^{0^{ud}}(4\,I_9(1^{uu}1^{dd}0^{ud}0^{ud})\\
&+\ 2\,I_9(1^{uu}0^{ud}1^{dd}0^{ud}) + 2\,I_9(1^{dd}0^{ud}1^{uu}0^{ud}) + 2\,I_{12}(1^{uu}0^{ud}1^{dd}0^{ud}) + 4\,I_{12}(1^{dd}1^{uu}0^{ud}0^{ud})\\
&+\ 3\,I_{12}(0^{ud}1^{uu}1^{dd}0^{ud})) + \alpha_1^{1^{dd}}(2\,I_9(1^{dd}0^{ud}1^{uu}1^{dd}) + I_{12}(0^{ud}1^{uu}1^{dd}1^{dd}))\\
&+\ \alpha_2^{1^{uu}0^{ud}}(4\,I_{13}(1^{uu}0^{ud}1^{dd}1^{uu}0^{ud}) + 2\,I_{13}(0^{ud}1^{uu}1^{dd}1^{uu}0^{ud}) + 4\,I_{14}(1^{uu}0^{ud}1^{dd}0^{ud}1^{uu})\\
&+\ 4\,I_{14}(1^{dd}0^{ud}1^{uu}0^{ud}1^{uu}) + 4\,I_{15}(1^{uu}0^{ud}1^{dd}1^{uu}0^{ud}) + 4\,I_{15}(1^{uu}0^{ud}1^{dd}0^{ud}1^{uu})\\
&+\ 2\,I_{15}(0^{ud}1^{uu}1^{dd}1^{uu}0^{ud})) + \alpha_2^{0^{ud}0^{ud}}(2\,I_{13}(1^{dd}1^{uu}0^{ud}0^{ud}0^{ud}) + 2\,I_{13}(0^{ud}1^{uu}1^{dd}0^{ud}0^{ud})\\
&+\ 8\,I_{14}(1^{uu}1^{dd}0^{ud}0^{ud}0^{ud}) + 4\,I_{14}(1^{uu}0^{ud}1^{dd}0^{ud}0^{ud}) + 4\,I_{14}(1^{dd}0^{ud}1^{uu}0^{ud}0^{ud})\\
&+\ 8\,I_{15}(1^{uu}1^{dd}0^{ud}0^{ud}0^{ud}) + 4\,I_{15}(1^{dd}1^{uu}0^{ud}0^{ud}0^{ud}) + 2\,I_{15}(0^{ud}1^{uu}1^{dd}0^{ud}0^{ud})\\
&+\ 4\,I_{15}(1^{dd}0^{ud}1^{uu}0^{ud}0^{ud}) + 4\,I_{15}(0^{ud}1^{dd}1^{uu}0^{ud}0^{ud})) + \alpha_2^{1^{uu}1^{dd}}(2\,I_{13}(0^{ud}1^{uu}1^{dd}1^{uu}1^{dd})\\
&+\ 4\,I_{15}(1^{dd}0^{ud}1^{uu}1^{dd}1^{uu})) + \alpha_2^{1^{dd}0^{ud}}(4\,I_{13}(1^{dd}1^{uu}0^{ud}0^{ud}1^{dd}) + 2\,I_{15}(1^{dd}0^{ud}1^{uu}1^{dd}0^{ud})\\
&+\ 4\,I_{15}(0^{ud}1^{dd}1^{uu}1^{dd}0^{ud})) + 4\,\alpha_3^{1^{uu}1^{uu}0^{ud}} I_{18}(0^{ud}1^{uu}1^{dd}1^{uu}1^{uu}0^{ud})\\
&+\ \alpha_3^{1^{uu}0^{ud}0^{ud}}(4\,I_{16}(1^{uu}0^{ud}1^{dd}0^{ud}1^{uu}0^{ud}) + 4\,I_{16}(1^{uu}0^{ud}1^{dd}0^{ud}0^{ud}1^{uu}) + 8\,I_{18}(1^{dd}1^{uu}0^{ud}0^{ud}1^{uu}0^{ud})\\
&+\ 4\,I_{18}(0^{ud}1^{uu}1^{dd}0^{ud}1^{uu}0^{ud}) + 4\,I_{18}(1^{uu}0^{ud}1^{dd}0^{ud}1^{uu}0^{ud})) + \alpha_3^{1^{uu}0^{ud}1^{dd}}(4\,I_{16}(1^{dd}0^{ud}1^{uu}0^{ud}1^{dd}1^{uu})\\
&+\ 4\,I_{18}(0^{ud}1^{dd}1^{uu}1^{uu}0^{ud}1^{dd}) + 4\,I_{18}(1^{dd}0^{ud}1^{uu}0^{ud}1^{uu}1^{dd})) + \alpha_3^{0^{ud}0^{ud}0^{ud}}(8\,I_{16}(1^{uu}1^{dd}0^{ud}0^{ud}0^{ud}0^{ud})\\
&+\ 4\,I_{16}(1^{dd}0^{ud}1^{uu}0^{ud}0^{ud}0^{ud}) + 4\,I_{18}(1^{uu}1^{dd}0^{ud}0^{ud}0^{ud}0^{ud}) + 4\,I_{18}(1^{dd}0^{ud}1^{uu}0^{ud}0^{ud}0^{ud}))\\
&+\ 4\,\alpha_3^{0^{ud}0^{ud}1^{dd}} I_{18}(0^{ud}1^{dd}1^{uu}0^{ud}0^{ud}1^{dd})
\end{aligned}\tag{A13}$$

$$\begin{aligned}
\alpha_3^{0^{ud}0^{ud}0^{ud}} &= \lambda + \alpha_1^{1^{uu}}(3\,I_9(0^{ud}0^{ud}0^{ud}1^{uu}) + 6\,I_{12}(0^{ud}0^{ud}0^{ud}1^{uu})) + \alpha_1^{0^{ud}}(6\,I_9(0^{ud}0^{ud}0^{ud}0^{ud})\\
&+\ 9\,I_{12}(0^{ud}0^{ud}0^{ud}0^{ud})) + \alpha_1^{1^{dd}}(3\,I_9(0^{ud}0^{ud}0^{ud}1^{dd}) + 3\,I_{12}(0^{ud}0^{ud}0^{ud}1^{dd}))\\
&+\ 6\,\alpha_2^{1^{uu}1^{uu}} I_{14}(0^{ud}0^{ud}0^{ud}1^{uu}1^{uu}) + \alpha_2^{1^{uu}0^{ud}}(6\,I_{13}(0^{ud}0^{ud}0^{ud}1^{uu}0^{ud}) + 24\,I_{14}(0^{ud}0^{ud}0^{ud}1^{uu}0^{ud})\\
&+\ 12\,I_{15}(0^{ud}0^{ud}0^{ud}1^{uu}0^{ud}) + 12\,I_{15}(0^{ud}0^{ud}0^{ud}0^{ud}1^{uu})) + \alpha_2^{0^{ud}0^{ud}}(6\,I_{13}(0^{ud}0^{ud}0^{ud}0^{ud}0^{ud})\\
&+\ 18\,I_{14}(0^{ud}0^{ud}0^{ud}0^{ud}0^{ud}) + 18\,I_{15}(0^{ud}0^{ud}0^{ud}0^{ud}0^{ud})) + \alpha_2^{1^{uu}1^{dd}}(6\,I_{13}(0^{ud}0^{ud}0^{ud}1^{uu}1^{dd})
\end{aligned}$$

$$
\begin{aligned}
&+\; 12\,I_{14}(0^{ud}0^{ud}0^{ud}1^{uu}1^{dd}) + 12\,I_{15}(0^{ud}0^{ud}0^{ud}1^{dd}1^{uu}) + 6\,I_{15}(0^{ud}0^{ud}0^{ud}1^{uu}1^{dd}))\\
&+\; \alpha_2^{1^{dd}0^{ud}}(12\,I_{14}(0^{ud}0^{ud}0^{ud}0^{ud}1^{dd}) + 6\,I_{15}(0^{ud}0^{ud}0^{ud}0^{ud}1^{dd}) + 6\,I_{15}(0^{ud}0^{ud}0^{ud}1^{dd}0^{ud}))\\
&+\; \alpha_3^{1^{uu}1^{uu}1^{dd}}(6\,I_{16}(0^{ud}0^{ud}0^{ud}1^{uu}1^{dd}1^{uu}) + 12\,I_{18}(0^{ud}0^{ud}0^{ud}1^{uu}1^{uu}1^{dd}))\\
&+\; \alpha_3^{1^{uu}0^{ud}0^{ud}}(6\,I_{16}(0^{ud}0^{ud}0^{ud}0^{ud}1^{uu}0^{ud}) + 12\,I_{16}(0^{ud}0^{ud}0^{ud}0^{ud}0^{ud}1^{uu}) + 12\,I_{18}(0^{ud}0^{ud}0^{ud}0^{ud}1^{uu}0^{ud})\\
&+\; 12\,I_{18}(0^{ud}0^{ud}0^{ud}1^{uu}0^{ud}0^{ud})) + \alpha_3^{1^{uu}0^{ud}1^{dd}}(12\,I_{16}(0^{ud}0^{ud}0^{ud}0^{ud}1^{uu}1^{dd})\\
&+\; 12\,I_{16}(0^{ud}0^{ud}0^{ud}1^{uu}1^{dd}0^{ud}) + 12\,I_{16}(0^{ud}0^{ud}0^{ud}1^{dd}0^{ud}1^{uu}) + 12\,I_{18}(0^{ud}0^{ud}0^{ud}1^{dd}1^{uu}0^{ud})\\
&+\; 12\,I_{18}(0^{ud}0^{ud}0^{ud}0^{ud}1^{uu}1^{dd})) + \alpha_3^{0^{ud}0^{ud}0^{ud}}(12\,I_{16}(0^{ud}0^{ud}0^{ud}0^{ud}0^{ud}0^{ud})\\
&+\; 12\,I_{16}(0^{ud}0^{ud}0^{ud}0^{ud}0^{ud}0^{ud}))
\end{aligned}
\tag{A14}
$$

$$
\begin{aligned}
\alpha_3^{0^{ud}0^{ud}1^{dd}} &=\; \lambda + \alpha_1^{1^{uu}}(I_9(0^{ud}0^{ud}1^{dd}1^{uu}) + 6\,I_{12}(0^{ud}0^{ud}1^{dd}1^{uu})) + \alpha_1^{0^{ud}}(4\,I_9(1^{dd}0^{ud}0^{ud}0^{ud}) + 2\,I_9(0^{ud}0^{ud}1^{dd}0^{ud})\\
&+\; 6\,I_{12}(1^{dd}0^{ud}0^{ud}0^{ud}) + 6\,I_{12}(0^{ud}0^{ud}1^{dd}0^{ud})) + \alpha_1^{1^{dd}}(4\,I_9(1^{dd}0^{ud}0^{ud}1^{dd}) + I_9(0^{ud}0^{ud}1^{dd}1^{dd}))\\
&+\; 6\,\alpha_2^{1^{uu}1^{uu}}I_{14}(0^{ud}0^{ud}1^{dd}1^{uu}1^{uu}) + \alpha_2^{1^{uu}0^{ud}}(12\,I_{13}(0^{ud}0^{ud}1^{dd}1^{uu}0^{ud}) + 24\,I_{14}(1^{dd}0^{ud}0^{ud}0^{ud}1^{uu})\\
&+\; 12\,I_{14}(0^{ud}0^{ud}1^{dd}1^{uu}0^{ud}) + 6\,I_{15}(0^{ud}0^{ud}1^{dd}1^{uu}0^{ud}) + 6\,I_{15}(0^{ud}0^{ud}1^{dd}0^{ud}1^{uu}))\\
&+\; \alpha_2^{0^{ud}0^{ud}}(6\,I_{13}(1^{dd}0^{ud}0^{ud}0^{ud}0^{ud}) + 24\,I_{14}(1^{dd}0^{ud}0^{ud}0^{ud}0^{ud}) + 6\,I_{14}(0^{ud}0^{ud}1^{dd}0^{ud}0^{ud})\\
&+\; 12\,I_{15}(1^{dd}0^{ud}0^{ud}0^{ud}0^{ud}) + 12\,I_{15}(0^{ud}1^{dd}0^{ud}0^{ud}0^{ud}) + 6\,I_{15}(0^{ud}0^{ud}1^{dd}0^{ud}0^{ud}))\\
&+\; \alpha_2^{1^{uu}1^{dd}}(12\,I_{15}(1^{dd}0^{ud}0^{ud}1^{dd}1^{uu}) + 6\,I_{15}(0^{ud}0^{ud}1^{dd}1^{dd}1^{uu})) + 12\,\alpha_2^{1^{dd}0^{ud}}I_{15}(0^{ud}1^{dd}0^{ud}1^{dd}0^{ud})\\
&+\; 12\,\alpha_3^{1^{uu}1^{uu}0^{ud}}I_{18}(0^{ud}0^{ud}1^{dd}1^{uu}1^{uu}0^{ud}) + 6\,\alpha_3^{1^{uu}1^{uu}1^{dd}}I_{16}(0^{ud}0^{ud}1^{dd}1^{uu}1^{dd}1^{uu})\\
&+\; \alpha_3^{1^{uu}0^{ud}0^{ud}}(6\,I_{16}(0^{ud}0^{ud}1^{dd}0^{ud}1^{uu}0^{ud}) + 12\,I_{16}(0^{ud}0^{ud}1^{dd}0^{ud}0^{ud}1^{uu}) + 12\,I_{18}(0^{ud}1^{dd}0^{ud}1^{uu}0^{ud}0^{ud})\\
&+\; 12\,I_{18}(0^{ud}0^{ud}1^{dd}0^{ud}1^{uu}0^{ud}) + 24\,I_{18}(1^{dd}0^{ud}0^{ud}0^{ud}1^{uu}0^{ud})) + 24\,\alpha_3^{1^{uu}0^{ud}1^{dd}}I_{16}(1^{dd}0^{ud}0^{ud}0^{ud}1^{dd}1^{uu})\\
&+\; \alpha_3^{0^{ud}0^{ud}0^{ud}}(24\,I_{16}(1^{dd}0^{ud}0^{ud}0^{ud}0^{ud}0^{ud}) + 12\,I_{18}(0^{ud}1^{dd}0^{ud}0^{ud}0^{ud}0^{ud}))
\end{aligned}
\tag{A15}
$$